

Dental and Alveolar Arch Widths in Normal Occlusion and Class II Division 1 Malocclusion

Jamal K. Mahmud BDS, MSc.⁽¹⁾

Anas K. Hamdy BDS, MSc.⁽²⁾

Muhammed I. Hazem BDS, MSc.⁽³⁾

Key words

Normal occlusion ,
malocclusion.

Abstract

This study was to evaluate and compare dental arch and alveolar widths of patients with Class II division one malocclusion with group of untreated normal occlusion subjects. This study was performed using measurement on dental casts of 30 normal occlusion and 30 Class II division 1 malocclusion of Iraqi sample aged 15-22 years. T-test was applied for comparing the groups , the finding of this study indicated that maxillary interpremolar width, maxillary canine, maxillary premolar and maxillary first molar and mandibular premolar and molar alveolar widths were found significantly narrower in the Class II division 1 group when compared with the normal occlusion sample ($P<.001$). The maxillary and mandibular intermolar and mandibular intercanine alveolar widths were statistically significantly larger in the Class II division 1 group ($P<.01$) . Maxillary molar teeth in subjects with Class II division 1 malocclusion tend to incline to the buccal to compensate the insufficient alveolar base therefore rapid maxillary expansion rather than slow maxillary expansion should be considered during treatment of Class II division 1 patients. The results of our study suggested that transverse discrepancy in Class II division one malocclusion originated from upper posterior teeth and maxillary alveolar base.

Introduction

Class II malocclusion is reported as the most frequently seen skeletal disharmony in orthodontic population. Investigators have recommended strongly the early detection of all Classes of malocclusion. Furthermore, they endorse preventive and interceptive orthodontics and dentofacial orthopedics for young patients to avoid, or at least to minimize the occurrence of malocclusion at the adult stage¹.

The size and shape of the arches have considerable implications in orthodontic diagnosis and treatment planning, affecting the space available, dental esthetics, and stability of the dentition². The growth changes of arch widths in normal occlusion subjects and a comparison of arch widths in normal occlusion and different malocclusion samples have been studied extensively³⁻⁹. Most of the studies in the literature compare dental arch widths of Class II patients with the normal occlusion samples. Some of them indicate that absolute arch widths of children with malocclusion did not differ appreciably from those with normal occlusion^{3,4,6,8,9}.

(1) lecturer, Department of Preventive, Orthodontics and Pedodontics, College of Dentistry, University of Tikrit.

(2) Assistant lecturer Department of Preventive, Orthodontics and Pedodontics, College of Dentistry, University of Tikrit.

(3) lecturer, Department of Periodontolog, College of Dentistry, University of Tikrit.

However, in other studies, statistically significant differences were determined in dental and alveolar width measurements of Class II patients^{3,9}. Sayin and Turkkahraman³ compared the arch and alveolar widths of patients with Class II division 1 malocclusion and subjects with Class I ideal occlusion in the permanent dentition. They indicated that mandibular intercanine widths were significantly larger in the Class II division 1 group, although maxillary intermolar widths were larger in the normal occlusion sample. Staley et al⁴ stated that patients with Class II, division 1 malocclusion had narrower maxillary intercanine, intermolar, and alveolar widths. Buschang et al⁶ had evaluated the differences in dental arch morphology among untreated adult females with Class I, Class II, division 1, and Class II, division 2 malocclusions and reported that Class II, division 1 females had the longest and narrowest arches. Tollaro et al¹⁰ found that patients with Class II malocclusions had a significantly narrower maxillary area during the mixed dentition phase. Bishara et al¹¹ reported that transverse discrepancy in Class II division 1 subjects did not appear to be self-corrected from the deciduous to the permanent dentitions. Similarly, Baccetti et al¹² reported that transverse interarch discrepancy was evident in the deciduous dentition and persisted into the mixed dentition. This study compares arch and alveolar widths of patients with Class II division 1 malocclusion and subjects with Class I ideal occlusion in the permanent dentition.

Materials and Method

This study was performed on dental casts of 30 normal occlusion and 30 Class II division 1 malocclusion subjects who were attending Tikrit University, College of Dentistry, Department of Orthodontics and different healthy centers in Saladin city.

Normal Occlusion Sample

Dental casts were taken from 30 adult subjects (13 male and 17 female) with normal occlusion, which met the following

criteria:³ (1) Class I canine and molar relationship with minor or no crowding, normal growth and development, well-aligned upper and lower dental arches; (2) all teeth present except third molars; (3) good facial symmetry determined clinically; (4) no significant medical history; (5) no history of trauma; and (6) no previous orthodontic or prosthodontic treatment, maxillofacial or plastic surgery.

Malocclusion Sample

A sample of 30 subjects (16 male and 14 female) with Class II division 1 malocclusion were selected from patient records. The inclusion criteria used to select Class II division 1 samples were³ (1) bilateral Class II molar relationship in centric occlusion with the distobuccal cusp tip of the maxillary first molar within one mm (anterior or posterior) from the buccal groove of the mandibular first molar and protrusive maxillary incisors; (2) all teeth present except third molars; (3) no significant medical history; and (4) no history of trauma, and no previous orthodontic, prosthodontic treatment, maxillofacial or plastic surgery. Twelve width measurements were performed on the dental casts of each subject⁹. The arch width measurements were recorded from each subject's dental casts by one examiner, using a dial caliper and recording the data to the nearest 0.1 mm. These dental and alveolar arch width measurements are shown in Table 1, and Figure 1 and 2. Four weeks after the first measurements, 20 dental casts were selected randomly and remeasured. A paired samples t-test was applied to the measurements. The difference between the first and second measurements was statistically insignificant. Independent-samples t-test was applied for comparison of the groups. All statistical analyses were performed using the Statistical Package for Social Sciences for Windows (SPSS) software package (version 10.1, SPSS Inc, Chicago, Ill).

Results

Four weeks after the first measurements, 20 randomly selected dental casts were remeasured. A paired-samples t-test was

applied to the measurements. The difference between the first and second measurements of the 20 casts was statistically insignificant. The method error was calculated by using Dahlberg's formula. Values varied from 0.360 to 0.882 and were within acceptable limits (Table 2). Descriptive statistics (mean, standard deviation, and minimum and maximum) and statistical comparisons of dental and alveolar width measurements for dental casts in two groups (normal occlusion and Class II division 1 malocclusion) are shown in Table 3. According to independent samples t-test, statistically significant differences were found in maxillary and mandibular dental arch and alveolar width dimensions between Class II division 1 malocclusion and normal occlusion samples. Statistical comparisons of the two groups showed no significant differences in maxillary intercanine (UC-C), mandibular interpremolar (LP-P) and mandibular canine alveolar width (LAC-C) measurements ($P > .05$). Statistically significant differences were found in nine of the 12 measurements. The maxillary interpremolar width ($P < .001$); maxillary canine ($P < .001$), premolar ($P < .001$), and molar ($P < .05$) alveolar widths; and mandibular premolar ($P, .001$) and molar ($P < .05$) alveolar widths were significantly narrower in the Class II division 1 group when compared with the normal occlusion sample. The upper ($P < .01$) and lower ($P < .01$) intermolar and lower intercanine ($P < .001$) widths were statistically significantly larger in the Class II division 1 group (Table 3).

Discussion

Information regarding maxillary arch dimensions in human populations is important to clinicians in orthodontics, prosthodontics, and oral surgery. It also is of interest to anthropologists and other students of human oral biology¹³. A survey of arch size could help the clinician in choosing correctly shaped stock impression trays for prosthodontic treatment. In addition to the selection of stock trays, the sizes of artificial teeth and the overall form of the artificial dental

arch at the wax trial stage are amenable to modification by the dental surgeon in orthodontic treatment¹⁴. This study was carried out to compare the dental arch and alveolar base widths of Class II division 1 malocclusion groups with the untreated normal occlusion sample. Width measurements described in this article will help clinicians diagnose and plan the treatment of patients with Class II division 1 malocclusions samples. Investigators who studied growth changes in the transverse arch width found that molar and canine arch widths did not change after age 13 in female subjects and age 16 in male subjects¹⁵⁻¹⁹. The minimum ages of the subjects measured in this study were chosen on the basis of these previous studies. Therefore, we assumed that the arch widths of the subjects studied were fully developed. Clinicians have speculated that nasal obstruction, finger habits, tongue thrusting, low tongue position, and abnormal swallowing and sucking behaviors were reasons for narrower maxillary dental arch widths in Class II division 1 malocclusions compared with a normal occlusion sample. Staley et al⁴ stated that the maxillary dental arch as a whole is narrower in adults with Class II division 1 malocclusion than it is in adults with normal occlusion. When we compare the dental and alveolar arch widths of Class II division 1 malocclusion samples with the normal occlusion samples, statistically significant lower values were found in most of the upper arch widths in Class II division 1 patients. Intercanine widths were investigated in a few of the previous studies, and conflicting results were found. These differences may be due to the age or severity of malocclusion of the subjects examined. Frohlich²⁰ compared intercanine widths of both arches from children with Class II malocclusion with data collected by Moorrees¹⁹ from children with normal occlusion and found that absolute arch widths of the Class II children did not differ appreciably from those of children with normal occlusion. Staley et al⁴ reported that subjects with normal occlusion had larger maxillary canine widths than the malocclusion subjects, but no differences were found in mandibular

canine widths. Bishara et al¹¹ studied the growth trends in maxillary and mandibular dental arch widths and lengths in persons with Class II division 1 malocclusions and normal subjects and reported no differences in maxillary and mandibular canine width measurements between the groups. In contrast with the others, Sayin and Turkkahraman³ found that mandibular intercanine widths were significantly larger in the Class II division 1 group than in the Class I group, whereas no significant differences were found among maxillary intercanine width measurements. In accordance with Sayin and Turkkahraman³, the results of this study showed that the maxillary intercanine width difference was similar in Class I and Class II division 1 groups, and the mandibular intercanine width was significantly larger in the Class II division 1 sample. Intermolar widths were the most commonly evaluated measurement in previous studies. Frohlich²⁰ found no difference in molar widths between normal and Class II subjects. Bishara et al¹¹ found no significant differences in the female comparisons of the intermolar width measurements. Staley et al⁴ reported that subjects with normal occlusion had larger maxillary molar widths and intermolar width differences than subjects with malocclusion. While evaluating alveolar widths, they reported that maxillary alveolar widths and mandibular alveolar widths of the males were larger in the Class I group. Again, alveolar width differences were found to be greater in the Class I group. They suggested that palatal movement of maxillary posterior teeth in Class II patients was needed to compensate for the increased overjet and to have good posterior interdigitation. Tollaro et al¹⁰ compared arch widths of 60 Class II, division 1 patients (26 males and 34 females) with 70 Class I subjects (25 males and 35 females) in the mixed dentition. Class II, division 1 subjects were grouped according to the presence of the posterior transverse interarch discrepancy (PTID). They reported that Class II, division 1 patients with PTID had narrower maxillary intermolar widths than Class II, division 1 patients without PTID

and Class I subjects. Mandibular intermolar widths did not differ between the three groups. They also suggested that Class II patients with PTID needed a preliminary expansion of the maxillary arch. In this study, molar reference points were taken from Staley et al⁴, who measured the widths between the mesiobuccal cusp tips of the maxillary first molars and the buccal grooves of the lower first molars. Because, in normal centric occlusion, the mesiobuccal cusp tips of the maxillary molars are positioned near the buccal grooves of the mandibular molars.

In this study we suggested that the narrow widths of the dental arch in Class II division 1 patients appeared to be caused by palatally tipped teeth and also by narrower bony bases of the dental arch. The results showed that transverse discrepancy in Class II division 1 patients originated from upper posterior teeth and not from the maxillary alveolar base. However, in contrast with previous studies, these findings indicated that the upper alveolar intermolar width was narrower and upper and lower intermolar widths were larger in patients with Class II division 1 malocclusion when compared with the normal occlusion sample. Therefore, we concluded that subjects with Class II division 1 malocclusions tend to have the maxillary molar teeth inclined to the buccal to compensate for the insufficient alveolar base. For that reason, rapid maxillary expansion rather than slow expansion may be considered before or during the treatment of a Class II division 1 patient, in agreement with Staley et al⁴ and Sayin and Turkkahraman³.

Of the four main categories in Angle's classifications of malocclusions, the Class II division 2 type of discrepancy occurs the least often. Obtaining data on Class II division 2 patients has always been challenging because of the low prevalence rates. For that reason, little data was found in the literature related to the alveolar widths of this malocclusion. In one of them, Walkow and Peck²¹ indicated that mandibular canine width was significantly less in Class II division 2 deepbite patients and suggested that the extreme deep bite may inhibit anterior development of the

mandibular dentoalveolar segment. In a cross-sectional study, Buschang et al⁶ found that Class II division 2 patients showed smaller mandibular intercanine and intermolar widths than the Class I and II division 1 patients. Buschang et al⁶ reported that in an adult female sample, the maxillary intermolar width of Class II cases was smaller than in the Class I subjects. Within the class II malocclusion group, the maxillary width was smaller in the Class II division 1 group than in the Class II division 2 subjects. Lux et al²² found dental arch widths of Class II division 2 cases took a position between the Class II division 1 cases and the Class I control groups. Moorrees et al¹⁹ used serial dental casts of untreated Class II malocclusions to compare arch dimensions of Class II division 1 and Class II division 2 subgroups. Compared with dental cast measurements from a control reference population, the Class II division 2 dental casts had greater than average maxillary and mandibular intercanine distances with the intermolar distances normally distributed. The findings of this study indicated that almost all the upper dental and alveolar width measurements were narrower in patients with Class II division 1 malocclusion when compared with the normal occlusion sample. In addition, the mandibular dental width measurements were larger in the Class II division 1 group. Subjects with Class II division 1 malocclusion tend to have the maxillary teeth inclined to the lingual and mandibular teeth inclined to the buccal direction because of the restriction of maxillary growth and development. Therefore, rapid maxillary expansion may be considered before or during the treatment of a Class II division 1 patient.

2. Based on previous studies on relapse, it is generally agreed that post orthodontic occlusal stability is enhanced through maintenance of the original inter-canine width. Again no significant differences found comparing upper and lower intermolar widths (M-M) in all groups, this come in accordance with Frhlich⁽³¹⁾, Nojima et al⁽³²⁾, Kook⁽³³⁾ and Susan and Elham⁽¹¹⁾, while disagree with other studies^(34,35) in which the M-M width is class II division 1 is significantly smaller

than class II division 2 and class I normal occlusion. Lastly comparing arch length (I-CC, I-MM) of all groups there were no significant differences except that of upper and lower arch length in class II division 1 is significantly larger than that of class II division 2 and normal occlusion this is in accordance to the finding of Susan and Elham in Jordan population⁽¹¹⁾, this may be attributed to proclination of anterior teeth. Whereas, comparing with class I only upper anterior arch length of class I normal occlusion is significantly smaller than class II division 1 as well as the lower I-MM. As a suggestion to further studies, it is of interest to relate dental arch width to the prevalence of respiratory disease. Epidemiologic surveys have shown that the prevalence of asthma in Iraq is relatively higher than in other nearby countries like Turkey, Lebanon, Jordan. Moreover, and was much higher than France, Italy, Sweden and Spain⁽³⁶⁾. Accordingly, further study is needed to investigate the association between the prevalence of respiratory disease and arch widths in our population.

Conclusions

-Maxillary interpremolar width, all maxillary alveolar widths, and mandibular premolar and molar alveolar widths were significantly narrower in the Class II division 1 group when compared with the normal occlusion sample.

-Upper alveolar intermolar width was narrower, and upper and lower intermolar width was greater in patients with Class II division 1 malocclusion when compared with the normal occlusion sample.

-Maxillary molar teeth in subjects with Class II division 1 malocclusions tend to incline buccally to compensate for the insufficient alveolar base. For that reason, rapid maxillary expansion rather than slow expansion may be considered before or during the treatment of a Class II division 1 patient.

Table(1):- Maxillary and mandibular dental and alveolar width measurements used in the study

1. Maxillary intercanine width (UC-C): the distance between the cusp tips of the right and left canines or the center of the wear facets in cases of attrition.
2. Maxillary interpremolar width (UP-P): the distance between the cusp tips of the right and left first premolars.
3. Maxillary intermolar width (UM-M): the distance between the mesiobuccal cusp tips of the right and left first molars.
4. Mandibular intercanine width (LC-C): the distance between the cusp tips of the right and left mandibular canines.
5. Mandibular interpremolar width (LP-P): the distance between the cusp tips of the right and left mandibular first premolars.
6. Mandibular intermolar width (LM-M): between the most gingival extensions of the buccal grooves on the first molars or, when the grooves had no distinct terminus on the buccal surface, between points on the grooves located at the middle of the buccal surfaces.
7. Maxillary canine alveolar width (UAC-C): the distance between two points at the mucogingival junctions above the cusp tips of the maxillary right and left canines.
8. Maxillary premolar alveolar width (UAP-P): the distance between two points at the mucogingival junctions above the interdental contact point of the maxillary first and second premolars.
9. Maxillary molar alveolar width (UAM-M): the distance between two points at the mucogingival junctions above the mesiobuccal cusp tips of the maxillary first molars
10. Mandibular canine alveolar width (LAC-C): the projection of UAC-C point in the lower jaw
11. Mandibular premolar alveolar width (LAP-P): the projection of UAP-P point in the lower jaw
12. Mandibular molar alveolar width (LAM-M): the projection of UAM-M point in the lower jaw




Fig.(1):- Mandibular dental cast measurements modified from (Sayin and Turkkahraman).




Fig.(2) :-Maxillary dental cast measurements modified from(Sayin and Turkkahraman)

Table(2):- Error of the method

Transverse measurment	Dahlberg's calculation	Reliability coefficient
UC-C	0.882	0.993
UP-P	0.567	0.912
UM-M	0.786	0.983
LC-C	0.654	0.917
LP-P	0.756	0.903
LM-M	0.360	0.982
UAC-C	0.534	0.943
UAP-P	0.456	0.936
UAM-M	0.365	0.948
LAC-C	0.452	0.915
LAP-P	0.654	0.920
LAM-M	0.423	0.936

Table(3):- Descriptive statistics and statistical comparisons of dental and alveolar widths of normal occlusion and class II division 1 malocclusion samples ^a

	Normal occlusion				Class II division 1 malocclusion				Normal occlusion vs class II div 1
	Mean	SD	Min	Max	Mean	SD	Min	Max	
UC-C	33.4	2.2	28.1	39.5	32.2	2.2	26.1	37.6	NS
UP-P	41.1	3.1	33.2	49.8	36.1	3.1	30.2	47.4	***
UM-M	51.3	2.6	45.0	60.1	53.3	2.6	47.0	63.3	**
LC-C	24.9	4.2	21.1	34.2	26.9	4.2	23.4	39.5	***
LP-P	33.6	1.6	28.3	40.4	34.6	1.6	29.3	41.5	NS
LM-M	46.2	1.7	40.3	52.1	48.3	1.7	41.3	54.2	**
UAC-C	38.5	3.6	32.6	45.9	35.5	3.6	30.5	42.9	***
UAP-P	48.9	4.7	40.8	58.7	44.3	4.7	37.8	55.0	***
UAM-M	59.4	2.2	55.6	65.3	56.4	2.2	53.4	62.3	*
LAC-C	33.8	3.4	30.2	41.0	32.7	3.4	29.2	40.2	NS
LAP-P	46.6	2.3	41.2	52.3	40.1	2.3	36.3	46.3	***
LAM-M	59.1	2.8	52.4	68.4	56.1	2.8	50.4	68.3	***

^a a SD indicates standard deviation; Min, minimum; Max, maximum; NS, not significant.

* P< 0.05, ** P< 0.01, *** P< 0.001.

References

- 1- El-Mangoury NH, Mostafa YA. Epidemiologic panorama of dental occlusion. Angle Orthod. 1990;60:207–214.
- 2- Lee RT. Arch width and form: a review. Am J Orthod Dentofacial Orthop. 1999;115:305–313.
- 3-Sayin MO, Turkkahraman H. Comparison of dental arch and alveolar widths of patients with Class II division 1 malocclusion and subjects with Class I ideal occlusion. Angle Orthod.2004;74:356–360.
- 4- Staley RN, Stuntz WR, Peterson LC. A comparison of arch widths in adults with normal occlusion and adults with Class II division 1 malocclusion. Am J Orthod. 1985;88:163–169.
- 5- Enlow DH, Hans MG. Essentials of Facial Growth. Philadelphia,Pa: WB Saunders; 1996:1–280.
- 6- Buschang PH, Stroud J, Alexander RG. Differences in dental arch morphology among adult females with untreated Class I and Class II malocclusion. Eur J Orthod. 1994;16:47–52.
- 7- Moorrees CFA, Gron AM, Le Bret LML, Yen PKJ, Frohlich FJ. Growth studies of the dentition: a review. Am J Orthod.1969;55:600–616.
- 8- Walkow TM, Peck S. Dental arch width in Class II division 2 deep-bite malocclusion. Am J Orthod Dentofacial Orthop.2002;122:608–613.
- 9-Uysal T, Memili B, Usumez S, Sari Z. Dental and alveolar arch widths in normal occlusion, Class II

division 1 and Class II division 2. *Angle Orthod.* 2005;75(6):756–762.

10-Tollaro I, Baccetti T, Franchi L, Tanesescu CD. Role of posterior transverse interarch discrepancy in Class II, division 1 malocclusion during the mixed dentition phase. *Am J Orthod Dentofacial Orthop.* 1996;110:417–422.

11-Bishara SE, Bayati P, Jakobsen JR. Longitudinal comparisons of dental arch changes in normal and untreated Class II, division 1 subjects and their clinical implications. *Am J Orthod Dentofacial Orthop.* 1996;110:483–489.

12-Baccetti T, Franchi L, McNamara JA Jr, Tollaro I. Early dentofacial features of Class II malocclusion: a longitudinal study from the deciduous through the mixed dentition. *Am J Orthod Dentofacial Orthop.* 1997;11:502–509.

13-Younes SAES. Maxillary arch dimensions in Saudi and Egypt population sample. *Am J Orthod Dentofacial Orthop.* 1984;85:83–88.

14-Braun S, Hnat WP, Fender DE, Legan HL. The form of the human dental arch. *Angle Orthod.* 1998;68:29–36.

15-Knott VB. Size and form of the dental arches in children with good occlusion studied longitudinally from age 9 years to late adolescence. *Am J Phys Anthropol.* 1961;19:263–284.

16- Knott VB. Longitudinal study of dental arch widths at four stages of dentition. *Angle Orthod.* 1972;42:387–394.

17- DeKock WH. Dental arch depth and width studied longitudinally from 12 years of age to adulthood. *Am J Orthod.* 1972;62:56–66.

18- Sillman JH. Dimensional changes of the dental arches: longitudinal study from birth to 25 years. *Am J Orthod.* 1964;50:824–842.

19- Moorrees CFA. *The Dentition of the Growing Child.* Cambridge, Mass: Harvard University Press; 1959:87–110.

20- Frohlich FJ. A longitudinal study of untreated Class II type malocclusion. *Trans Eur Orthod Soc.* 1961;37:137–159.

21- Walkow TM, Peck S. Dental arch width in Class II division 2 deep-bite malocclusion. *Am J Orthod Dentofacial Orthop.* 2002;122:608–613.

22- Lux CJ, Conrath C, Burden D, Komposch G. Dental arch widths and mandibular-maxillary base widths in Class II malocclusions between early mixed and permanent dentitions. *Angle Orthod.* 2003;73:674–685.